elaborierte analyse t	Konfigura fréquent	ations-Frequ	enz-Analyse onfiguratio		K-LIST FOR OBSERVED CONFIGURATIO Strichliste der Anzahl beobachteter Konfig porée no. aux observations	urationen (Beobachtung	gswert %);	%); expecting value; Erwartungswert; expectation e	chi-square; Chi-Quadrat; ; chi carré ;	Standard	Fehlerwahrs	cheinlichkei Freiheitso	t~Halbierur grade;df~	T-HALF VALIDIT ngscheck ob gültig 4 -1 ; 2-1) BIP row-validity
\backslash	F dime	nsions of	4 configura	ations ·			1		T			 I	T	
nr.	<i>pattern</i> (Gf)	<i>of class</i> (Au)	ified catego (Aw)	<i>ories</i> (Am	stripe for each unit according to ob b) (no. RUN; RUN %)	served categories	Σ (o %)	e% = 6,25%	$\chi^2 = \Sigma (0.6, 25)^2 : 6, 25$	α<5%;	α < 1%	α < 5%	α < 1%	α; 1 ^{st/2} ~ 2 nd
01.	+	+	+	+						*7,81	**13,3	~	~	•
02.	+	+	+	-						*7,81	**13,3	~	~	
03.	+	+	-	+						*7,81	**13,3	~	~	
04.	+	+	-	-						*7,81	**13,3	~	~	
05.	+		+	+						*7,81	**13,3	~	~	
06.	+	-	+	-						*7,81	**13,3	~	~	
07.	+	-	-	+						*7,81	**13,3	~	~	
08.	+	-	-	-						*7,81	**13,3	~	~	
09.	-	+	+	+						*7,81	**13,3	~	~	
10.	-	+	+	-						*7,81	**13,3	~	~	
11.	-	+	-	+						*7,81	**13,3	~	~	
12.	-	+	-	-						*7,81	**13,3	~	~	
13.	·		+	+						*7,81	**13,3		~	
14.	-	-	+	-						*7,81	**13,3	~	~	
15.	-	-	-	+						*7,81	**13,3	~	~	
16.	-	-	-	-						*7,81	**13,3	~	~	
	F CHECK	K; 2x Halbier	rungs-Iteratio	n (BIP):	percentage configuration patterns 1s	t & 2nd half	Σ (0%)	e % = 25%	(o-25) ² : 25	*5%; 3 df	**1%; 3 df	*5%; 1df	*1%; 1df	1. Hälfte ~ 2. H 1 <u>e m ~ 2e</u> m
SHC 01.	+	· _ · · _ · · -					+	·· _ ·· _ ·· _ ·· _	······	+··-·-·	' ►	*3,84	**6.64	
SHC 02.	+	_			ļ į					~	~	*3.84	**6.64	
SHC 02.		+								~	~	*3,84	**6.64	
SHC 03.	-	•								~	~	*3.84	**6.64	
STU 04.	-	-								~ 0	~ Kurt-W. L			l

Appendix: algorhythm to: Elaborated configurationfrequency-analysis, e-KFA

Algorythm after a hectographed contribution in a seminary lesson on social cognitions and behaviour at psychological institute, university of the Saar, Saarbrücken, summer-semester, 1975 by Kurt-Wilhelm Laufs, ©, updated as English version 2014-11-02, 2015-02-26

At a first glance, e-KFA could remind Cochran's Q-sort, yet is not. KFA had been formulated by Krauth & Lienert about 1971 to typologize and to analyse by chi-square and binominal distribution. Critics on KFA (Konfiguration – Frequenz -

Analyse) had followed lexically (Clauss, G. & al., 1976: Wörterbuch der Psychologie. VEB Verlag Enzyklopädie, Leipzig. Pahl-Rugenstein, Köln, 1976), and described the problem to smaller or larger number of checked persons or items than about N ~ 40.

This numerical methodical KFA inherent problem really can be avoided, if instead of absolute numeri one calculated in percentages, so one could also analyse rather approprietly, both, smaller samples than N ~ 40, or larger samples than N \sim 40. Author's KFA elaboration shows examples, how to apply e-KFA in psychology, and psychological field research. and also as a practitioner's method, without any electrical computer, just calculations bv hand to combinations of hypotheses social in any and psychological field.

Do it yourself !

1st you define your most possible to observe terms, categories, or dimensions, according to valid theories &/or objective items.

2nd you take the amount (number) of categories to form plus-minus (yes/no

signatures answers or as combinatoric plus/minus) configuration-matrices. A two configuration-matrix (KF)four makes possible configurative combinations: (++; +-; -+; --), a three KF shows eight configurative combinations (+++; ++-; +-+; +--; -++; -+-; --+; ---) etc. Above algorythm sheed shows four configurations with sixteen possible of "yes" &/or "no" configurations of, combinatorically, etc. Why now KFA elaborated, behalf to calculate in 4-When percentages? configurations were by split half (bi-partation) analysed chi-square, after only significant values in row after

split half iteration were valid. thus a four-configuration must be equally significant at least after chi-square BIP controll in it's both parts divided in two configurations. The lowest significance in determines here the row significance of all a fourconfiguration row. Percentage calculation as

appropriate to social and psychological data (always in mind that "nasty" scaling problem and of objectivity), claims percentages for numbers of observed data and to expecting values of a distribution as inference model. social When data or psychological data in practice

or social fields occur, that phantasm of normal or binominal distributions can even more appropriate and more rapidly be calculated by inference of equal distribution. Thus: 100% of postulated expecting inference (e) be to twoconfigurational percentage number observed (o) data, as 100% by 4 (number of possible combinations) = 25% expectation value (e); for 3-configuration's observed percentages (o) and it's possible combinations 100% by 8 = 12,5% (e); four configuration's e = 6,25 %; (100%:16).When one will look for

 α errors of significancy in

one's statistical tables on chi square, degrees of freedom (df; FG) depend here on number of configurations: two-configurations make 2 - 1 = 1 df; three-config. show 3-1 = 2 df; and fourconfigurative calculations make subtract one from four and show three degrees of freedom at it's table value for significance.

This appropriate and rather quick method to calculate shows very satisfying approximations to much more complicated factor analyses and also can be applied to control rapidly factor analyses by hand calculation without electronic computers, and also efficiently can be applied in social fields, on park banches, and in practice, to bundle data and test it's significance, without scaling and interpretation problems those problems typical to factor analysis of communality and rotation. Nevertheless one could bundle e-KFA results again by a factor analysis. To rapid e-KFA percentages can be done inter-correlations to test reliability to it's percentage results in rows (types/factors) and columns (factorial categories, items, dimensions), and an arithmetical (or geometrical by f. exampl. Mosier nomogram, 1942, in: Lienert, G.A., 1970: Testtheorie. Beltz. Weinheim) mean coefficient

can describe consistency (as well as a communality) coefficient.

Literature: see above text, within and at author's WEB-site,

Terms:

Psychology, psycho-linguistics, mathematical psychology, structuralisme, algorhythm to elaborated configuration frequency analysis (e-KFA), percentage chi-square, equal distribution, benefit by method: smaller and larger samples than N = 40 to calculate, e-KFA is an approximation to factor analysis; e-KFA "types" (factors) can be further condensed by factor analysis, when taking percentage types as "variables"; also to evaluate Rorschach (Ro) systematics psycho-linguistically.



Author and Copyright, © 1975 ff: Kurt-Wilhelm Laufs, D.P. (Diplom-Psychologe, phil. & min. med. fac.), ev. KiR i.R., Zum Resthof 2, D-23996 Bobitz, 2014-11-02, 2014-11-03, 2014-11-05, 2014-11-27, 2014-11-28, update 2015-02-26,©

No e-mails, rather picture postcards, please!